

## Gene Delivery to the Blood-Brain Barrier

## Author: Louiza Bohn Thomsen

Laboratory for Neurobiology – Biomedicine, Department of Health Science and Technology, Aalborg University, Aalborg, Denmark

Drug- and gene delivery to the brain is highly restricted by the vascular barriers of

the brain, denoted by the blood-brain barrier (BBB) and the blood-cerebrospinal

fluid (CSF) barriers. Among these barriers, BBB is the main limiting factor. It is

composed by the brain capillary endothelial cells (BCECs). The BCECs barrier function is supported by astrocytes, pericytes and neurons to form the

blood-brain barrier. BCECs are very tightly connected to each other by tight junctions.

Apart from the essential substrates needed to nourish the brain, small and/or lipophilic

molecules are free to diffuse into the brain. However most pharmacologically

active drugs and gene fragments are too large to enter the brain. Various kinds of

drug-carriers have been constructed for delivery of large substances to the brain.

Such drug-carriers have to be able to successfully carry their cargo through the

BCECs and into the brain. For testing the ability of drug-carriers to deliver their

cargo into the brain, investigators have constructed different in vitro BBB models,

consisting of BCECs that express the main characteristics of the BBB in vivo.

In the first part of the thesis the ability of two drug-carriers, pullulanspermine and SPIOs, to mediate transfection of BCECs or transcellular transport through BCECs in vitro was studied.<br/>

Pullulan-spermine is a polymeric complex consisting of the

polysaccharide, pullulan and the polyamine, spermine. Pullulan-spermine formed a

cationic complex shown to be able to bind plasmid DNA electrostatically.

Pullulan-spermine was conjugated with plasmid DNA encoding a red fluorescent

protein, Hc-Red-1 C1, or human growth hormone 1 (hGH1). Pullulan-spermine

complexed with Hc-Red-1 C1 cDNA led to the formation of a red fluorescent signal in human brain microvascular endothelial cells (HBMECs). Furthermore,

pullulan-spermine complexed with hGH1 cDNA was not only able to transfect

 $\ensuremath{\mathsf{HBMECs}}$  but also led to secretion of the hGH1 into the culture media. Pullulanspermine-

cDNA complexes could transfect non-dividing cells although the rate of transgene cells was higher in dividing cells. This indicated that the DNA is

not only entering the cell nucleus under mitosis. Unfortunately, pullulan-spermine

complexes proved incapable of transfecting HBMECs in the presence of serum in

the growth media and additional studies are needed to enable its use for in vivo

transfection.<br><br>

Another potential drug-carrier, fluorescent iron oxide nanoparticles were

also shown to enter HBMECs upon incubation. These nanoparticles were also able

to pass though the HBMECs forming a BBB in a static in vitro BBB model. Furthermore, their passage was increased by the aid of an external magnetic field

created by placing the cell culture plates with the SPIOs on a plate magnet. Two

vitality tests showed no significant change in BCEC vitality after addition of Non-viral delivery strategies into and across the brain capillary endothelial cells xiv

SPIOs or by dragging the nanoparticles through the BCECs in the presence

of the

external electric field.<br><br>

The results of the drug-carrier studies indicate that it is possible to deliver plasmid cDNA into BCECs and transfect these cells leading to their

secretion of encoded protein into the extracellular space. Moreover, SPIOs are potentially

potent carriers of attachable molecules trough cultured BCECs in vitro, which may

have high potential for drug-delivery to the brain in vivo.<br><br>

In the second part of the thesis, two in vitro BBB models, a static and a

dynamic model was investigated and compared. The static model consisting of

microporous membrane inserts in which immortalized BCECs is cultured. The model induces many characteristics of the BBB in vivo, but lacks the

tiahtness induction factor of shear stress. Different experiments were performed with

this static model to monitor BBB integrity. Barrier formation by the BCECs was

monitored by measuring transendothelial electric resistance (TEER) and the BCFC

monolayer was stained positive for zonula occludens 1 (ZO-1) a tight junction

protein. It was mainly found that the tightness of the BCECs was strengthened by

contact co-culture of the BCECs with astrocytes and addition of hydrocortisone to

the media. The dynamic in vitro BBB model however, did not lead to any reliable

results in this study and further investigation of barrier formation in this model was not pursued. In consequence a comparison between the static and dynamic

in vitro models was not possible, but it could be concluded that the static model

seems to

be the most reliable model

## Laboratory for Neurobiology

Gene Delivery to the Blood-Brain Barrier: Non-Viral Gene Carriers and in vitro **Blood-Brain Barrier Models** 



PhD thesis by Louiza Bohn Thomsen



## **River Publishers Series in open**

ISBN: 9788792982148 e-ISBN: 9788792982155 Available From: December 2012 Price: € 40.00

**KEYWORDS:** astrocyte-conditioned media



www.riverpublishers.com marketing@riverpublishers.com